



Office: 323-442-5860

Cell: 248-721-3740

Email: kevincollon@gmail.com

SHOULDER HEMIARTHROPLASTY PROCEDURE REHABILITATION PROTOCOL

	RANGE OF MOTION	IMMOBILIZER	EXERCISES
PHASE I 0-4 weeks	Limit ER to passive 45° to protect subscap repair PROM FE to 135, Abduction to 90, ABER 45, ABIR 30 No active IR nor extension until 6 weeks	0-2 weeks: Worn at all times (day and night) Off for gentle exercise only 2-4 weeks: Worn daytime only	0-2 weeks: Grip strengthening, pendulum exercises Elbow/wrist/hand ROM at home 2-4 weeks: Begin cuff, deltoid isometrics; limit ER 45° passive Supine, upright PROM, modalities
PHASE II 4-12 weeks	Increase as tolerated to full Begin active assisted/active internal rotation and extension as tolerated after 6 weeks	None	6-8 weeks: Begin light resisted ER, forward flexion and abduction, active IR. 8-12 weeks: Begin resisted internal rotation, extension and scapular retraction
PHASE III 12-24 weeks	Progress to full motion without discomfort	None	Advance strengthening as tolerated Closed chain scapular rehab and functional rotator cuff strengthening; focus on anterior deltoid and teres Maximize subscapular stabilization